

Gas-solid coexistence of adhesive spheres

P. Charbonneau and D. Frenkel*

*FOM Institute for Atomic and Molecular Physics,
Kruislaan 407, 1098 SJ Amsterdam, The Netherlands*

(Dated: February 3, 2008)

The Baxter limit of adhesive spheres¹ is often deemed an unphysical model for liquids, due to the thermodynamic metastability of its disordered phases with respect to crystal formation^{2,3}. Here we suggest conditions under which the disordered phases are actually stable with respect to the solid for narrow attractive interactions down to the Baxter limit. Furthermore, using basic free energy considerations we estimate the necessary condition for the gas-liquid critical point to be stable with respect to crystallization⁴. Possible experimental and simulation realizations are briefly discussed.

For what follows, it is convenient to consider the original Baxter adhesive sphere model, i.e. a square-well fluid with well depth ϵ and well width δ , in hard-core diameter σ units, which we set equal to unity for convenience. Except where explicitly mentioned, the results are for three-dimensional systems. For adhesive spheres, the stickiness τ behaves as a temperature-like quantity, which is defined as

$$\tau^{-1} \equiv 12\delta e^{\epsilon/k_B T}, \quad (1)$$

where $k_B T$ is the product of Boltzmann's constant with temperature. The equivalent quantity for square-well potentials with finite width⁵ is

$$\tau^{-1} = 4 \left((1 + \delta)^3 - 1 \right) \left(e^{\epsilon/k_B T} - 1 \right). \quad (2)$$

The second virial coefficient can then be expressed as $B_{2v}/B_{2v}^{HS} = 1 - \frac{1}{4\tau}$, where $B_{2v}^{HS} = 2\pi/3$ is the hard-sphere value.

The critical point of sticky spheres (and, to a good approximation, *all* fluids with a narrow, isotropic attraction⁶) occurs at $\tau_c \approx 0.11$ ⁴. How about freezing? At low temperatures, the density of the gas coexisting with the solid is very low, $\rho_g \ll 1$, while at high temperatures, the freezing density is that of hard spheres. Typically, the freezing curve moves rather abruptly, but continuously, from one limit to the other. We estimate the temperature and corresponding τ where this happens and compare it with the known τ_c ⁴.

To estimate the crossover temperature where the density of the vapor coexisting with the solid starts to increase, we make use of the fact that the chemical potential per particle μ is the same in the different coexisting phases. For the fluid, we use an ideal gas (we check this approximation later), while we treat the crystal as a collection of uncorrelated cells each with free volume v_f . Although this last approximation is rather crude compared to what is usually used in similar systems⁹, for narrow square-well potentials at low temperatures, it is




FIG. 1: Gas-solid coexistence line from Eq. 4 for square-well interaction with $Z = 12$ and $\delta = 0.25, 0.15$, and 0.03 , from top to bottom. Simulation results for the first two cases are included for comparison^{7,8}.




FIG. 2: Sketch of the three-dimensional Baxter phase diagram following ref.⁴ for the isotropic gas-liquid binodal (dashed line) and this work for the gas-solid coexistence curves (solid lines). The gas-solid coexistence is found at high τ for $Z > 6$ and at very low τ for $Z < 6$, while it follows Eq. 7 for $Z = 6$. Gas-crystal coexistence curves for $\rho \gtrsim 0.1$ and gas-liquid coexistence for $\tau < 0.1$ are drawn for schematic purposes only.

not unreasonable. This gives

$$\begin{aligned} \frac{\Delta\mu}{k_B T} &= \frac{\Delta e}{k_B T} - \frac{\Delta s}{k_B} + \frac{p\Delta v}{k_B T} \\ &\approx -\frac{Z\epsilon}{2k_B T} - \ln \rho_g v_f - 1, \end{aligned} \quad (3)$$

where s is the entropy, v is the volume, e is the energy, and Z is the coordination number of the crystal. To obtain the last line, we also assume v to be much smaller in the crystal than in the gas, so $\Delta v \approx \rho_g^{-1}$. Note that when $\Delta\mu = 0$, $k_B T_{coex}/Z\epsilon$ is the only temperature term, thus changing the coordination number of the solid effectively rescales the coexistence temperature. In other words, to a first approximation, the gas coexistence curve

for a solid with coordination number Z_2 can be obtained from the known gas coexistence curve for a solid with coordination number Z_1 by multiplying T_{coex} by Z_2/Z_1 .

Following Sear², we take $v_f \approx (\frac{\delta}{2})^d$ for narrow square wells in d dimensions. Eq. 3 can then be used to calculate gas-solid phase coexistence directly,

$$\frac{k_B T_{coex}}{\epsilon} = \frac{-Z/2}{\ln \rho_g \left(\frac{\delta}{2}\right)^d + 1}. \quad (4)$$

This is compared to published numerical results of $Z = 12$ square-well phase diagrams in Fig. 1. At low ρ_g , towards the wider limit of narrow wells, there is a fairly good agreement between this crude treatment and the MC simulated phase diagrams. Results for $Z = 4$ also qualitatively agree with this picture¹⁰.

As a last consistency check, we examine in what regime corrections to $\Delta v \approx \rho_g^{-1}$ are small. The next order terms are the crystal's finite volume, which is of $\mathcal{O}(1)$, and the fluid's second-virial coefficient B_{2v} . Using Eq. 4 as a mean-field-like closure for ϵ/T_{coex} gives

$$\begin{aligned} \frac{B_{2v}^{coex}}{B_2^{HS}} &\approx 1 - 3\delta \left(e^{\epsilon/T_{coex}} - 1 \right) \\ &\approx 1 - 3\delta \left(\left(\frac{8}{e\delta^3 \rho_g} \right)^{2/Z} - 1 \right), \end{aligned} \quad (5)$$

which for $Z \geq 6$ leaves an $\mathcal{O}(\rho_g^{-2/Z})$ correction. At low densities the two contributing factors are both $\ll \rho_g^{-1}$ and have an opposite sign, so they amount to an error in T_{coex} of less than a few percents for narrow well widths when $\rho_g \lesssim 0.1$. This level of precision is sufficient for the sake of our argument.

Going to the Baxter limit, $\delta \rightarrow 0$, we then use the definition of stickiness from Eq. 1 to rewrite Eq. 4 as

$$1 = -\ln \left[2^{-d} \rho_g \delta^{d-Z/2} \left(\frac{1}{12\tau_{coex}} \right)^{Z/2} \right]. \quad (6)$$

For $Z > 2d$, $\delta^{d-Z/2}$ diverges, while at freezing the RHS of Eq. 6 should be $\mathcal{O}(1)$. If we fix the density of the gas at a small but finite value, it follows that $\tau_{coex} \rightarrow \infty$, i.e. the fluid freezes well above $\tau_c \approx 0.11$. Conversely, when $Z < 2d$, $\tau_{coex} = 0$, so the triple point also lies on the $\tau = 0$ axis. The interesting case is the marginal situation $Z = 2d$, where we obtain a non-trivial fixed crossover τ_{coex} . For $d = 3$, the crossover τ_{coex} increases

by less than 15% as δ varies from 0.25 down to the Baxter limit, which is small relative to the divergence obtained for $Z > 2d$. Moreover we expect τ_{coex} to lie in the same regime as τ_c . Indeed, in three dimensions Eq. 6 reduces to

$$\tau_{coex} = \frac{(e\rho_g)^{1/3}}{24}. \quad (7)$$

Using $\rho_g = 0.1$ as a reference point, we then obtain a crossover $\tau_{coex} \approx 0.03$, which is well below τ_c for the isotropic case, as shown in Fig. 2. The gas-liquid critical point is thus *stable* with respect to the crystal.

Simple cubic crystals, among others, and isostatic solids, fulfill the $Z = 2d$ condition for all d . On the other hand, monodisperse, isotropically adhesive spheres, have invariably $Z = 12$ crystal phases. However, if either the spheres are not monodisperse or if the interaction is patchy, a lower Z can be obtained. In the adhesive limit, even infinitesimally polydisperse solids eventually fractionate², reaching thermodynamic equilibrium, which then results in $Z \approx 12$. However, this is kinetically nearly impossible and therefore of little experimental relevance. Instead, an isostatic polydisperse metastable crystal would leave τ_c stable with respect to the gas-solid coexistence. For patchy adhesive spheres Z is precisely controlled by the number of patches. Though τ_c also varies with Z ¹¹, this effect is unlikely to be sufficient to bring it below τ_{coex} .

In conclusion, if one manages to reduce the bonding of the solid phase down to $Z \leq 2d$, the high density fluid remains stable with respect to the solid down to the Baxter limit. Based on rough, but reasonable estimates, the three-dimensional gas-liquid critical point would then be also stable with respect to crystallization. Furthermore, this scheme might be realizable both experimentally and through simulation.

Acknowledgments

PC would like to thank D.R. Reichman for insightful discussions. PC also acknowledges Marie-Curie IIF XXXXXXXXX funding. This work is part of the research program of the ‘‘Stichting voor Fundamenteel Onderzoek der Materie (FOM)’’, which is financially supported by the ‘‘Nederlandse organisatie voor Wetenschappelijk Onderzoek (NWO)’’.

* Electronic address: frenkel@amolf.nl

¹ R. J. Baxter, J. Chem. Phys. **49**, 2770 (1968).

² R. P. Sear, Mol. Phys. **96**, 1013 (1999).

³ G. Stell, J. Stat. Phys. **63**, 1203 (1991).

⁴ M. A. Miller and D. Frenkel, Phys. Rev. Lett. **90**, 135702 (2003).

⁵ M. A. Miller and D. Frenkel, J. Phys.: Condens. Matter **16**, S4901 (2004).

⁶ M. G. Noro and D. Frenkel, J. Chem. Phys. **113**, 2941 (2000).

⁷ D. L. Pagan and J. D. Gunton, J. Chem. Phys. **122**, 184515 (2005).

- ⁸ H. Liu, S. Garde, and S. Kumar, J. Chem. Phys. **123**, 174505 (2005).
- ⁹ C. F. Tejero, A. Daanoun, H. N. W. Lekkerkerker, and M. Baus, Physical Review E **51**, 558 (1995).
- ¹⁰ Z. Zhang, A. S. Keys, T. Chen, and S. C. Glotzer, *Langmuir* **21**, 11547 (2005).
- ¹¹ E. Bianchi, J. Largo, P. Tartaglia, E. Zaccarelli, and F. Sciortino, Phys. Rev. Lett. **97**, 168301 (2006).